

ORIENTABILITY OF LINEAR WEINGARTEN SURFACES, SPACELIKE CMC-1 SURFACES AND MAXIMAL SURFACES

MASATOSHI KOKUBU AND MASAOKI UMEHARA

ABSTRACT. We prove several topological properties of linear Weingarten surfaces of Bryant type, as wave fronts in hyperbolic 3-space. For example, we show the orientability of such surfaces, and also co-orientability when they are not flat. Moreover, we show an explicit formula of the non-holomorphic hyperbolic Gauss map via another hyperbolic Gauss map which is holomorphic. Using this, we show the orientability and co-orientability of CMC-1 faces (i.e., constant mean curvature one surfaces with admissible singular points) in de Sitter 3-space. (CMC-1 faces might not be wave fronts in general, but belong to a class of linear Weingarten surfaces with singular points.) Since both linear Weingarten fronts and CMC-1 faces may have singular points, orientability and co-orientability are both nontrivial properties. Furthermore, we show that the zig-zag representation of the fundamental group of a linear Weingarten surface of Bryant type is trivial. We also remark on some properties of non-orientable maximal surfaces in Lorentz-Minkowski 3-space, comparing the corresponding properties of CMC-1 faces in de Sitter 3-space.

1. INTRODUCTION

Let M^2 and N^3 be C^∞ manifolds of dimension 2 and of dimension 3, respectively. The projectified cotangent bundle $P(T^*N^3)$ has a canonical contact structure. A C^∞ map $f : M^2 \rightarrow N^3$ is called a *frontal* if f lifts to a Legendrian map L_f , i.e., a C^∞ map $L_f : M^2 \rightarrow P(T^*N^3)$ such that the image $dL_f(TM^2)$ of the tangent bundle TM^2 lies in the contact hyperplane field on $P(T^*N^3)$. Moreover, f is called a *wave front* or a *front* if it lifts to a Legendrian immersion L_f . Frontals (and therefore fronts) generalize immersions, as they allow for singular points. A frontal f is said to be *co-orientable* if its Legendrian lift L_f can lift up to a C^∞ map into the cotangent bundle T^*N^3 , otherwise it is said to be *non-co-orientable*. It should be remarked that, when N^3 is Riemannian, a front f is co-orientable if and only if there is a globally defined unit normal vector field ν along f . (In [21], ‘fronts’ were implicitly assumed to be co-orientable by definition, and fronts which are not necessarily co-orientable were distinguished as different, by calling them ‘p-fronts’. However, in this paper, a front may be either co-orientable or non-co-orientable and the term ‘p-front’ is not used.)

Date: July 20, 2010.

1991 *Mathematics Subject Classification.* Primary: 53A35; Secondary: 53C40, 53C42.

Key words and phrases. linear Weingarten, wave front, (co-)orientability, hyperbolic space, de Sitter space, zig-zag representation.

The first author was supported by Grant-in-Aid for Scientific Research (C) No. 18540096 from the Japan Society for the Promotion of Science. The second author was supported by Grant-in-Aid for Scientific Research (A) No. 19204005 from the Japan Society for the Promotion of Science.

In [14], Gálvez, Martínez and Milán gave a fundamental framework for the theory of linear Weingarten surfaces of Bryant type in hyperbolic 3-space H^3 . We shall investigate such surfaces in H^3 and in de Sitter 3-space S_1^3 in the category of (wave) fronts which admit certain kinds of singular points. We remark that orientability and co-orientability are not equivalent for fronts, though they are equivalent for immersed surfaces. We prove that non-flat linear Weingarten fronts of Bryant type in H^3 and in S_1^3 are co-orientable and orientable. Flat fronts, in contrast, are orientable but are not necessarily co-orientable (see [21]). For a co-orientable front $f : M^2 \rightarrow H^3$, a representation of the fundamental group

$$\sigma_f : \pi_1(M^2) \rightarrow \mathbf{Z}$$

called the ‘zig-zag representation’ is induced (see Section 3.2), which is invariant under continuous deformations of fronts. We shall show that the zig-zag representation of any linear Weingarten front of Bryant type is trivial (see Theorem 3.9). Moreover, we give an explicit formula for the hyperbolic Gauss maps of these surfaces.

On the other hand, we shall also study CMC-1 faces (defined by Fujimori [7]) in S_1^3 . CMC-1 faces are constant mean curvature one surfaces on their regular sets, and are frontals (see Corollary 4.3), but not fronts in general. In fact, each limiting tangent plane at a singular point contains a lightlike vector. They also belong to a class of linear Weingarten surfaces of Bryant type and satisfy an Osserman-type inequality (cf. [7] and [9]) under a certain global assumption. As an application of an explicit formula for hyperbolic Gauss maps, we prove the orientability and the co-orientability of CMC-1 faces in S_1^3 , which is, in fact, deeper than the above corresponding assertion for linear Weingarten fronts, because CMC-1 fronts in S_1^3 admit only isolated singular points (see Corollary 3.4).

It should be remarked that CMC-1 faces in S_1^3 have quite similar properties to maxfaces (i.e. maximal surfaces in Lorentz-Minkowski 3-space $\mathbf{R}_1^3$ with admissible singular points defined in [28]). In Section 5 (this section is a joint work with Shoichi Fujimori and Kotaro Yamada), using the same method as in Section 4, we show the existence of globally defined real analytic normal vector field on a given maxface in $\mathbf{R}_1^3$.

Throughout this paper, M^2 denotes a smooth and connected 2-dimensional manifold or a Riemann surface; $\mathbf{R}_1^4$ denotes the Lorentz-Minkowski space of dimension 4, with the Lorentz metric

$$\langle (x_0, x_1, x_2, x_3), (y_0, y_1, y_2, y_3) \rangle = -x_0y_0 + x_1y_1 + x_2y_2 + x_3y_3,$$

and the hyperbolic 3-space H^3 and the de Sitter 3-space S_1^3 are defined by

$$H^3(= H_+^3) = \left\{ (x_0, x_1, x_2, x_3) \in \mathbf{R}_1^4; -x_0^2 + x_1^2 + x_2^2 + x_3^2 = -1, x_0 > 0 \right\},$$

$$S_1^3 = \left\{ (x_0, x_1, x_2, x_3) \in \mathbf{R}_1^4; -x_0^2 + x_1^2 + x_2^2 + x_3^2 = 1 \right\},$$

respectively. H^3 is a simply-connected Riemannian 3-manifold with constant sectional curvature -1 , and S_1^3 is a simply-connected Lorentzian 3-manifold with constant sectional curvature 1 . We occasionally consider

$$H_-^3 := \left\{ (x_0, x_1, x_2, x_3) \in \mathbf{R}_1^4; -x_0^2 + x_1^2 + x_2^2 + x_3^2 = -1, x_0 < 0 \right\}$$

isometric to H_+^3 .

2. LINEAR WEINGARTEN SURFACES OF BRYANT TYPE AS WAVE FRONTS

Let $f : M^2 \rightarrow H^3(\subset \mathbf{R}_1^4)$ be a front in hyperbolic 3-space H^3 .

If f is co-orientable, then there exists a C^∞ map $\nu : M^2 \rightarrow S_1^3$ satisfying $\langle f, \nu \rangle = 0$ and $\langle df, \nu \rangle = 0$, namely,

$$L_f = (f, \nu) : M^3 \rightarrow T_1 H^3 \subset H^3 \times S_1^3$$

is the Legendrian lift of f . (Here, we identify the unit tangent bundle $T_1 H^3$ with the unit cotangent bundle $T_1^* H^3$.) We call the C^∞ map ν a unit normal field of f . Interchanging their roles, ν can be considered as a spacelike front in S_1^3 with f as a unit normal field. Here, a front in S_1^3 is said to be *spacelike* if the plane in $T_{\nu(p)} S_1^3$ orthogonal to the normal vector $f(p) \in H^3$ is spacelike at each point p . (Conversely, for a given spacelike co-orientable front in S_1^3 , its unit normal field is a front which lies in the upper half or the lower half of the hyperboloid $H_+^3 \cup H_-^3$ of two sheets in $\mathbf{R}_1^4$.) For each real number δ , the pair

$$f_\delta = (\cosh \delta)f + (\sinh \delta)\nu, \quad \nu_\delta = (\cosh \delta)\nu + (\sinh \delta)f$$

gives a new front f_δ called a *parallel front* of f whose unit normal field is ν_δ .

From now on, we consider a front $f : M^2 \rightarrow H^3$ which may not be co-orientable. The following fact implies that even when the front f has a singular point p , taking a co-orientable neighborhood U of p . Then we can consider a parallel front

$$f_{U,\delta} := (\cosh \delta)f + (\sinh \delta)\nu \quad (\delta \in \mathbf{R})$$

on U , where ν is a unit normal vector field of f on U . For a suitable $\delta \in \mathbf{R}$, we may assume that $f_{U,\delta}$ is an immersion at p .

Fact 2.1. *Let $f : M^2 \rightarrow H^3$ be a front. Then for each $p \in M^2$, taking a co-orientable neighborhood U of p , the parallel fronts $f_{U,\delta}$ are immersions at p except for at most two values $\delta \in \mathbf{R}$.*

First, we begin with the case $f : M^2 \rightarrow H^3$ is an immersion, which is more restrictive than the case f is a front.

Definition 2.2. An immersion $f : M^2 \rightarrow H^3$ is said to be *horospherical linear Weingarten* if the mean curvature H (with respect to the normal field ν) and the Gaussian curvature K satisfy the relation $a(H-1) + bK = 0$ for some real constants a and b such that $a^2 + b^2 \neq 0$.

Remark 2.3. A horospherical linear Weingarten immersion $f : M^2 \rightarrow H^3$ with $a = 0$ is just a flat immersion, hence M^2 is orientable (see [21, Theorem B]). On the other hand, a horospherical linear Weingarten surface with $a \neq 0$ is also orientable by the following reasons:

- (1) Suppose moreover that f is minimal, i.e., $H = 0$. The horospherical linear Weingarten condition implies that the Gaussian curvature K is constant. Thus f is a totally geodesic immersion (see [5]).
- (2) Suppose f is not minimal. The opposite unit normal field $-\nu$ does not give a horospherical linear Weingarten immersion, that is, $(-H) - 1$ is never proportional to K . Thus it cannot happen that M^2 is non-orientable.

Remark 2.4. The parallel surfaces f_δ of a horospherical linear Weingarten immersion f are also horospherical linear Weingarten on the set R_{f_δ} of regular points of f_δ (see [19]). In particular, parallel surfaces of a flat surface are also flat on R_{f_δ} , which is a well known fact.

Fact 2.1 and Remark 2.4 enable us to give the following definition.

Definition 2.5. A front $f : M^2 \rightarrow H^3$ is called a *horospherical linear Weingarten front* if for each $p \in M^2$, there exist a co-orientable neighborhood U of p and $\delta_0 \in \mathbf{R}$ such that the parallel front f_{U, δ_0} is a horospherical linear Weingarten immersion on U . (Indeed, the parallel front $f_{U, \delta}$ is a horospherical linear Weingarten immersion at p with at most two exceptional values of $\delta \in \mathbf{R}$.)

We can give two exceptional examples of horospherical linear Weingarten fronts:

- (1) a hyperbolic line as a degeneration of a parallel surface of a hyperbolic cylinder,
- (2) a single point as a degeneration of a parallel surface of a geodesic sphere.

These two examples have no regular points in their domains of definition. Conversely, any horospherical linear Weingarten fronts not having regular points are locally congruent to one of these two examples (see the appendix). So we exclude them from our study in this paper. In other words, we will assume (except in the appendix) that *a horospherical linear Weingarten front has regular points*.

Note that a horospherical linear Weingarten front of zero Gaussian curvature is just a flat p-front defined in [21]. (See also [22] and [20].) For horospherical linear Weingarten fronts which are not flat, the following assertion holds:

Theorem 2.6. *Let $f : M^2 \rightarrow H^3$ be a non-flat horospherical linear Weingarten front. Then f is co-orientable. Moreover, there exists a unique ratio $[a : b]$ of real constants ($a^2 + b^2 \neq 0$) such that the parallel front f_δ satisfies*

$$(2.1) \quad a(H_\delta - 1) + b_\delta K_\delta = 0, \quad b_\delta := be^{2\delta} + \frac{a(e^{2\delta} - 1)}{2}$$

on its regular set R_{f_δ} , where H_δ and K_δ are the mean curvature and the Gaussian curvature of f_δ on R_{f_δ} , respectively.

Proof. Fix a regular point $p \in M^2$ arbitrarily. Let the equation $a(H - 1) + bK = 0$ ($a^2 + b^2 \neq 0$) hold for $f|_U$ with the unit normal field $\nu^{(p)}$ on a neighborhood U of p . Then the equation (2.1) holds on $U \cap R_{f_\delta}$ because of [19, (2.18)].

Take another point q close to p so that there exist a neighborhood U_q of q and $\delta_0 \in \mathbf{R}$ satisfying $U_q \cap U \neq \emptyset$ and f_{U_q, δ_0} is a horospherical linear Weingarten immersion with respect to the unit normal field $\nu^{(q)}$. Then $f_{U_q \cap U, \delta_0}$ is horospherical linear Weingarten with respect both to $\nu^{(p)}$ and to $\nu^{(q)}$. It follows from Remark 2.3 that $\nu^{(q)}$ coincides with $\nu^{(p)}$ on $U_q \cap U$. Moreover, since the equation (2.1) holds on $U_q \cap U$, it must hold on U_q .

Continuing this argument, we have the collection $\{\nu^{(q)} \mid q \in M^2\}$ which then fixes a global unit normal field ν . Hence the front f is co-orientable.

Finally, we note that the argument above implies the equation (2.1) holds on each $U_q \cap R_{f_\delta}$, therefore on the regular set R_{f_δ} . $\square$

Let $f : M^2 \rightarrow H^3$ be a horospherical linear Weingarten front which satisfies the relation $a(H - 1) + bK = 0$ for some real constants a and b ($a^2 + b^2 \neq 0$). Let $\nu : M^2 \rightarrow S_1^3$ denote the unit normal field of f (assuming the co-orientability whenever f is a flat front). Then $f + \nu$ is a map from M^2 to the lightcone Λ^3 in $\mathbf{R}_1^4$. Here, the lightcone Λ^3 is, by definition,

$$\Lambda^3 = \left\{ (x_0, x_1, x_2, x_3) \in \mathbf{R}_1^4; -x_0^2 + x_1^2 + x_2^2 + x_3^2 = 0 \right\}.$$

Except for the case where $a + 2b = 0$, the map $f + \nu$ induces a pseudometric of constant Gaussian curvature $\varepsilon = a/(a + 2b)$ (cf. [14], [19]). The equation $a(H - 1) + bK = 0$ can be rewritten as $2\varepsilon(H - 1) + (1 - \varepsilon)K = 0$.

Let $W_\varepsilon(M^2)$ denote the set of horospherical linear Weingarten fronts from M^2 to H^3 satisfying

$$\begin{cases} 2\varepsilon(H - 1) + (1 - \varepsilon)K = 0 & (\text{if } \varepsilon \in \mathbf{R}) \\ 2(H - 1) - K = 0 & (\text{if } \varepsilon = \infty). \end{cases}$$

- (Flat fronts) A front $f \in W_0(M^2)$ has zero Gaussian curvature, that is, $W_0(M^2)$ is the set of flat fronts defined on M^2 in H^3 . (In this case, $a = 0$.) The parallel fronts of a co-orientable flat front f also belong to $W_0(M^2)$. Fundamental properties of flat fronts are given in [13], [21] and [22].
- (Hyperbolic type) A front $f \in W_\varepsilon(M^2)$ is called a linear Weingarten front of *hyperbolic type* if $\varepsilon > 0$ (i.e., $a/(a + 2b) > 0$). The parallel fronts of f are also in the same class $\bigcup_{\varepsilon > 0} W_\varepsilon(M^2)$. (cf. [3], [27], etc., for the case $\varepsilon = 1$.) For each $f \in \bigcup_{\varepsilon > 0} W_\varepsilon(M^2)$, there exists a unique $\delta \in \mathbf{R}$ such that f_δ is a CMC-1 (constant mean curvature one) front in H^3 . In this case, the unit normal vector field $\nu_\delta : M^2 \rightarrow S_1^3$ is HMC-1 (harmonic-mean curvature one), that is, the mean of reciprocals of principal curvatures equals one. (See Corollary 3.3 for a related result.)
- (de Sitter type) A front $f \in W_\varepsilon(M^2)$ is called a linear Weingarten front of *de Sitter type* if $\varepsilon < 0$ (i.e., $a/(a + 2b) < 0$). The parallel fronts of f are also in the same class $\bigcup_{\varepsilon < 0} W_\varepsilon(M^2)$. For each $f \in \bigcup_{\varepsilon < 0} W_\varepsilon(M^2)$, there exists a unique $\delta \in \mathbf{R}$ such that f_δ is HMC-1 (see [19]). In this case, ν_δ gives the CMC-1 front in S_1^3 . (See Corollary 3.4 for a related result.)
- (Horo-flat fronts) A front $f \in W_\infty(M^2)$ is said to be *horo-flat* (cf. [16]). (In this case, $a + 2b = 0$.) If $f \in W_\infty(M^2)$, one of the hyperbolic Gauss maps (see Section 3) of f degenerates. The parallel fronts of f also belong to $W_\infty(M^2)$. Fundamental properties of horo-flat fronts are given in [16] and [26]. It should be remarked that there exist non-real analytic horo-flat surfaces (see [2, Examples 2.1 and 2.2]).

Remark 2.7. Our terminology ‘horospherical linear Weingarten’ comes from the following reasons: The horospheres in H^3 satisfy $K = H - 1 = 0$, and are elements in $W_1(M^2)$, $W_0(M^2)$, $W_{-1}(M^2)$ and $W_\infty(M^2)$ at the same time, as the degenerate cases. Recently, Izumiya et al. [15] proposed the horospherical geometry in H^3 , which includes all of the above classes $W_\varepsilon(M^2)$.

As pointed out at the beginning of this section, if $f : M^2 \rightarrow H^3$ is a co-orientable front, there exists a unit normal field ν defined globally on M^2 , which gives also a front $\nu : M^2 \rightarrow S_1^3$. For $f \in W_\varepsilon(M^2)$, one can verify that the mean curvature $\hat{H}$ and the Gaussian curvature $\hat{K}$ of $\nu : M^2 \rightarrow S_1^3$ satisfy

$$2\varepsilon(\hat{H} - 1) + (1 + \varepsilon)\hat{K} = 0.$$

This implies that flat fronts, horo-flat fronts, linear Weingarten fronts of hyperbolic type and of de Sitter type in H^3 correspond to those in S_1^3 , respectively.

Definition 2.8. Linear Weingarten fronts of hyperbolic type, of de Sitter type and flat fronts are all called *linear Weingarten fronts of Bryant type* (cf. [14]). In other

words, a linear Weingarten front of Bryant type is a non-horo-flat linear Weingarten front.

It should be remarked that Aledo and Espinar [1] recently classified complete linear Weingarten immersions of Bryant type with non-negative Gaussian curvature in S_1^3 . In this paper, almost all formulas are of linear Weingarten surfaces in H^3 , but one can find the several corresponding formulas of linear Weingarten surfaces in S_1^3 in [1].

3. ORIENTABILITY OF LINEAR WEINGARTEN FRONTS OF BRYANT TYPE AND AN EXPLICIT FORMULA FOR THE HYPERBOLIC GAUSS MAP G_*

We now concentrate on linear Weingarten fronts of Bryant type, i.e., fronts in $W_\varepsilon(M^2)$ ($\varepsilon \neq \infty$), which will be denoted by $f : M^2 \rightarrow H^3$ throughout this section.

3.1. A representation formula and singular points. The first, second and third fundamental forms are denoted by I , II and III , respectively. (In general, I and III are well-defined not only for immersions but also for fronts, and II is well-defined for co-orientable fronts.) The sum $I + III$ of the first and third fundamental forms of f is a positive definite metric on M^2 , because f is a front.

On the other hand, one can verify that the symmetric 2-tensor $\varepsilon I + (1 - \varepsilon)II$ for $f \in W_\varepsilon(M^2)$ ($\varepsilon \neq 0, \infty$) or for a co-orientable $f \in W_0(M^2)$ is definite on $M^2 \setminus S_f$ and vanishes on S_f , where S_f denotes the set of singular points of f . We now suppose that M^2 is orientable. Then there is a unique complex structure on the regular set $R_f := M^2 \setminus S_f$ so that $\varepsilon I + (1 - \varepsilon)II$ is Hermitian. Since this complex structure is common to the whole parallel family of f , and the singular set changes when taking a parallel surface (cf. Fact 2.1), it follows that *there is a unique complex structure on M^2 so that $\varepsilon I + (1 - \varepsilon)II$ is Hermitian.*

$\Lambda_+^3 \ni p \mapsto [p] \in \Lambda_+^3 / \sim$ denotes the canonical projection of the positive lightcone $\Lambda_+^3 = \{(x_0, x_1, x_2, x_3) \in \Lambda^3; x_0 > 0\}$ onto the set $\Lambda_+^3 / \sim$ of positively oriented lightlike lines. Note that $\Lambda_+^3 / \sim$ can be naturally identified with the Riemann sphere $S^2 \cong \mathbf{C} \cup \{\infty\}$. The *hyperbolic Gauss map* $G := [f + \nu]$ can be defined for $f \in W_\varepsilon(M^2)$ ($\varepsilon \neq 0$) and for a co-orientable flat front f .

From now until the end of this subsection, we assume that M^2 is orientable (though we will later know that this assumption is not necessary). And also, *throughout this section, all flat fronts considered are assumed to be co-orientable.* We have already shown that non-flat linear Weingarten surfaces of Bryant type are all co-orientable. On the other hand, even if $f : M^2 \rightarrow H^3$ is a non-co-orientable flat front, we can take a double covering $\pi : \hat{M}^2 \rightarrow M^2$ such that $f \circ \pi$ is co-orientable. So this assumption of co-orientability is not so restrictive.

The hyperbolic Gauss map $G : M^2 \rightarrow \mathbf{C} \cup \{\infty\} = S^2$ of f is a meromorphic function with respect to the complex structure mentioned above. On the other hand, the metric $d\sigma^2$ induced by $f + \nu : M^2 \rightarrow \Lambda_+^3$ is a Hermitian pseudometric on M^2 of constant curvature ε which may have isolated singular points of integral order (cf. [14], [19], the isolated singular points of $d\sigma^2$ in M^2 correspond to umbilical points of f). Then there exists a meromorphic function h on $\hat{M}^2$ satisfying

$$d\sigma^2 = \frac{4|dh|^2}{(1 + \varepsilon|h|^2)^2},$$

where $\tilde{M}^2$ is the universal covering of M^2 . (h is the so-called *developing map*.) Then the fundamental forms of f are written as follows (cf. [14], [19]):

$$(3.1) \quad I = \frac{(1-\varepsilon)^2}{4} d\sigma^2 + \frac{4|Q|^2}{d\sigma^2} + (1-\varepsilon)(Q + \bar{Q}),$$

$$(3.2) \quad II = \frac{\varepsilon^2 - 1}{4} d\sigma^2 + \frac{4|Q|^2}{d\sigma^2} - \varepsilon(Q + \bar{Q}),$$

$$(3.3) \quad III = \frac{(1+\varepsilon)^2}{4} d\sigma^2 - (1+\varepsilon)(Q + \bar{Q}) + \frac{4|Q|^2}{d\sigma^2},$$

where

$$(3.4) \quad Q := \frac{1}{2} (S(h) - S(G)) = \frac{1}{2} (\{h : z\} dz^2 - \{G : z\} dz^2).$$

The 2-differential Q is called the *Hopf differential* of f . (The Schwarzian derivative $S(G)$ as a locally defined meromorphic 2-differential is given by [21, (2.18)]. The difference $S(h) - S(G)$ does not depend on the choice of local complex coordinate z , see also [19, Lemma 3.7].) We also note that

$$\begin{aligned} \varepsilon I + (1-\varepsilon)II &= -\frac{(1-\varepsilon)^2}{4} d\sigma^2 + \frac{4|Q|^2}{d\sigma^2}, \\ I + III &= \frac{1+\varepsilon^2}{2} d\sigma^2 + \frac{8|Q|^2}{d\sigma^2} - 2\varepsilon(Q + \bar{Q}). \end{aligned}$$

The following assertion can be proved easily by a modification of [19, Proposition 3.5], which is a variant of the holomorphic representation formula in [14]:

Theorem 3.1. *Let G be a meromorphic function on a Riemann surface M^2 , and h a meromorphic function on the universal covering $\tilde{M}^2$ such that the pseudometric $d\sigma^2 := 4|dh|^2/(1+\varepsilon|h|^2)^2$ gives a single-valued symmetric tensor on M^2 . Suppose that*

$$(3.5) \quad \frac{1+\varepsilon^2}{2} d\sigma^2 + \frac{8|Q|^2}{d\sigma^2} - 2\varepsilon(Q + \bar{Q})$$

is positive definite, where $Q := (S(h) - S(G))/2 = (\{h : z\} dz^2 - \{G : z\} dz^2)/2$. Define $f = \mathcal{G}\mathcal{A}\mathcal{G}^$ and $\nu = \mathcal{G}\mathcal{B}\mathcal{G}^*$ by*

$$\mathcal{G} := i(G_h)^{-3/2} \begin{pmatrix} -GG_h & (GG_{hh}/2) - (G_h)^2 \\ -G_h & G_{hh}/2 \end{pmatrix},$$

where $G_h := dG/dh$, $G_{hh} := dG_h/dh$ and

$$\mathcal{A} := \begin{pmatrix} \frac{1+\varepsilon^2|h|^2}{1+\varepsilon|h|^2} & -\varepsilon\bar{h} \\ -\varepsilon h & 1+\varepsilon|h|^2 \end{pmatrix}, \quad \mathcal{B} := \begin{pmatrix} \frac{1-\varepsilon^2|h|^2}{1+\varepsilon|h|^2} & \varepsilon\bar{h} \\ \varepsilon h & -1-\varepsilon|h|^2 \end{pmatrix}.$$

Then $f : M^2 \rightarrow H^3 = SL(2, \mathbf{C})/SU(2)$ is a co-orientable linear Weingarten front of Bryant type in $W_\varepsilon(M^2)$, and $\nu : M^2 \rightarrow S_1^3 = SL(2, \mathbf{C})/SU(1, 1)$ is its unit normal. Moreover, G and (3.5) coincide with the hyperbolic Gauss map and the sum $I + III$ of the first and third fundamental forms of f , respectively. Conversely, any co-orientable linear Weingarten front of Bryant type in $W_\varepsilon(M^2)$, except for the horosphere, is given in this manner.

Remark 3.2. We give some additional explanation for Theorem 3.1:

(1) We often identify $\mathbf{R}_1^4$ with the set $\text{Herm}(2)$ of 2×2 Hermitian matrices by

$$(3.6) \quad X = (x_0, x_1, x_2, x_3) \leftrightarrow X = \sum_{k=0}^3 x_k e_k = \begin{pmatrix} x_0 + x_3 & x_1 + ix_2 \\ x_1 - ix_2 & x_0 - x_3 \end{pmatrix},$$

where $i = \sqrt{-1}$ and

$$e_0 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad e_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad e_2 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}, \quad e_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Since the Lorentzian inner product $\langle \cdot, \cdot \rangle$ is given by

$$\langle X, Y \rangle = -\frac{1}{2} \text{trace}(X e_2 {}^t Y e_2), \quad \langle X, X \rangle = -\det X,$$

H^3 and S_1^3 are given by

$$H^3 = \{X; X^* = X, \det X = 1, \text{trace} X > 0\},$$

$$S_1^3 = \{X; X^* = X, \det X = -1\},$$

respectively, where X^* denotes the transposed conjugate ${}^t \bar{X}$ to the matrix X . Hence, they are also represented by

$$H^3 = \{aa^*; a \in SL(2, \mathbf{C})\} = SL(2, \mathbf{C})/SU(2),$$

$$S_1^3 = \{ae_3a^*; a \in SL(2, \mathbf{C})\} = SL(2, \mathbf{C})/SU(1, 1).$$

(2) The choice of the developing map h of $d\sigma^2$ is not unique. However, the above expression of f does not depend on the choice of h . We prove this here:

Suppose that we have the following two expressions

$$d\sigma^2 = \frac{4|dh|^2}{(1 + \varepsilon|h|^2)^2} = \frac{4|d\tilde{h}|^2}{(1 + \varepsilon|\tilde{h}|^2)^2}.$$

By taking a parallel front, we may assume that f is an immersion on an open subset U of M^2 . Since f is real analytic, it is sufficient to show that $\tilde{f}$ associated to the pair $(G, \tilde{h})$ coincides with f . Since the first and second fundamental forms I and II can be written in terms of G , $d\sigma^2$ and Q as in (3.1) and (3.2), $\tilde{f}$ is congruent to f by the fundamental theorem of surface theory. Moreover, we can show that $f = \tilde{f}$ in this situation. In fact, $a = (a_{ij})_{i,j=1,2} \in SL(2, \mathbf{C})$ isometrically acts f as afa^* . Then the hyperbolic Gauss map G changed as $(a_{11}G + a_{12})/(a_{21}G + a_{22})$. Since $\tilde{f}$ and f have the same hyperbolic Gauss map G , which implies that $f = \tilde{f}$.

(3) $\mathcal{G}$ satisfies

$$(3.7) \quad \mathcal{G}^{-1}d\mathcal{G} = \begin{pmatrix} 0 & \theta \\ dh & 0 \end{pmatrix},$$

where $\theta = -\frac{1}{2}\{G : h\}dh$. Moreover, note that $Q = \theta dh$. (See [14] and [19] for details.)

Corollary 3.3. *Let $f \in W_\varepsilon(M^2)$ for $\varepsilon > 0$. Then there exists a unique real number δ such that $f_\delta : M^2 \rightarrow H^3$ is a CMC-1 front whose singular set S_f consists only of isolated points in M^2 .*

Proof. Without loss of generality, we may assume that f is not a horosphere. There exists a unique real number δ such that $f_\delta \in W_1(M^2)$. It follows from (3.1) with $\varepsilon = 1$ that the first fundamental form of f_δ is $I_{f_\delta} = 4|Q|^2/d\sigma^2$, which is positive definite except at the zeros of the holomorphic 2-differential Q . Since Q is holomorphic, f_δ is a front with only isolated singular points. $\square$

Similarly, we get the following:

Corollary 3.4. *Let $f \in W_\varepsilon(M^2)$ for $\varepsilon < 0$. Then there exists a unique real number δ such that the unit normal field ν_δ of f_δ gives a CMC-1 front $\nu_\delta : M^2 \rightarrow S_1^3$ whose singular set S_f consists only of isolated points in M^2 .*

Proof. Without loss of generality, we may assume that f is not a horosphere. There exists a unique real number δ such that $f_\delta \in W_{-1}(M^2)$, that is, f_δ has constant harmonic-mean curvature one (cf. [19]). At the same time, $\nu_\delta : M^2 \rightarrow S_1^3$ is a CMC-1 front. It follows from (3.1) and (3.3) with $\varepsilon = -1$ that the first fundamental form of ν_δ is $I_{\nu_\delta} = III_{f_\delta} = 4|Q|^2/d\sigma^2$. Since Q is holomorphic, ν_δ is a front with only isolated singular points. $\square$

Remark 3.5. A meromorphic function $G = z + iz^2$ and pseudometrics

$$d\sigma^2 = \frac{4|dh|^2}{(1+|h|^2)^2} \quad (\text{resp. } d\sigma^2 = \frac{4|dh|^2}{(1-|h|^2)^2}),$$

where $h = z + z^3$, induce CMC-1 fronts in H^3 (resp. in S_1^3). They have an isolated singular point at $z = 0$ because $S(h) - S(G) (= 2Q)$ vanishes at $z = 0$.

Remark 3.6. Let $f \in W_0(M^2)$ and ν its unit normal field. Then one can show that $f + \nu$ is a flat front in the lightcone in $\mathbf{R}_1^4$, whose singular set $S_{f+\nu}$ consists only of isolated points.

More generally, we can show the following propositions concerning singular points of linear Weingarten fronts of Bryant type.

Proposition 3.7. *Let $f : M^2 \rightarrow H^3$ be a linear Weingarten front of Bryant type written in terms of $(G, d\sigma^2 = 4|dh|^2/(1+\varepsilon|h|^2)^2)$ as in Theorem 3.1. Then the set of singular points equals*

$$S_f = \left\{ p \in M^2 ; \varepsilon I + (1-\varepsilon)II = 0 \right\} = \left\{ p \in M^2 ; \frac{4|Q|^2}{d\sigma^2} - \frac{(1-\varepsilon)^2}{4}d\sigma^2 = 0 \right\},$$

where Q is the Hopf differential (3.4). Moreover, a singular point $p \in S_f$ is nondegenerate if and only if

- (1) f is not CMC-1, and
- (2) $4\varepsilon h_z \bar{h} + (1+\varepsilon|h|^2)(\hat{\theta}_z/\hat{\theta} - h_{zz}/h_z) \neq 0$ holds at p , where $\theta = \hat{\theta}dz$ for a local coordinate z , and $h_z = dh/dz$, $h_{zz} = d^2h/dz^2$.

We now set

$$\Delta := \text{Im} \left[\frac{1}{\sqrt{1-\varepsilon}} \left\{ \frac{4\varepsilon h_z \bar{h}}{1+\varepsilon|h|^2} + \frac{\hat{\theta}_z}{\hat{\theta}} - \frac{h_{zz}}{h_z} \right\} \frac{1}{\sqrt{h_z \bar{\theta}}} \right],$$

where $\sqrt{1-\varepsilon}$ is an imaginary number when $1-\varepsilon < 0$.

Proposition 3.8. *For a nondegenerate singular point p of a linear Weingarten front $f : M^2 \rightarrow H^3$ of Bryant type, the germ of f at p is locally diffeomorphic to*

- (1) a cuspidal edge if and only if $\Delta \neq 0$ at p , or
- (2) a swallowtail if and only if $\Delta = 0$ at p and $\frac{d}{dt}\Big|_{t=0} \Delta \circ \gamma \neq 0$, where γ is a parametrization of S_f with $\gamma(0) = p$.

The proofs of the above propositions for the case of $\varepsilon = 0$ are given in [20]. The proofs for the case of $\varepsilon \neq 0$ can be followed by a quite similar argument to that in [20], though we need very lengthy calculations. We omit them here.

3.2. Zig-zag representation and orientability. For a given co-orientable front $f : M^2 \rightarrow H^3$, the zig-zag representation

$$\sigma_f : \pi_1(M^2) \rightarrow \mathbf{Z}$$

is induced (cf. [23], [24] and [25]), which is invariant under the deformation of f as a wave front, and

$$(3.8) \quad \sigma_f = \sigma_\nu$$

holds for the unit normal field $\nu : M^2 \rightarrow S_1^3$.

Theorem 3.9. *Let $f : M^2 \rightarrow H^3$ (resp. $f : M^2 \rightarrow S_1^3$) be a co-orientable linear Weingarten front of Bryant type. Then M^2 is orientable. Moreover, the zig-zag representation σ_f is trivial.*

Proof. It is sufficient to prove the assertion for a co-orientable linear Weingarten front $f : M^2 \rightarrow H^3$. (In fact, suppose that the assertion holds for linear Weingarten front of Bryant type in H^3 . Let $f : M^2 \rightarrow S_1^3$ be a co-orientable linear Weingarten front. Then its unit normal vector field $\nu : M^2 \rightarrow H^3$ is a co-orientable linear Weingarten front of Bryant type. Thus M^2 must be orientable. On the other hand, the zig-zag representation is trivial by (3.8).)

Suppose that $f : M^2 \rightarrow H^3$ is of hyperbolic type (resp. de Sitter type). By Corollary 3.3 (resp. Corollary 3.4), there exists $\delta \in \mathbf{R}$ such that f_δ (resp. ν_δ) is a CMC-1 front with only isolated singular points. Since f_δ (resp. ν_δ) is an immersion of non-vanishing mean curvature on $R_{f_\delta} = M^2 \setminus S_{f_\delta}$, the open submanifold R_{f_δ} is orientable. Since S_{f_δ} is discrete, we conclude that M^2 is also orientable. On the other hand, we can take a loop γ which represents a given element of $\pi_1(M^2)$ so that γ does not pass through S_{f_δ} . Since the zig-zag representation is trivial when the corresponding loop does not meet the singular set, $\sigma_{f_\delta}([\gamma]) = \sigma_{\nu_\delta}([\gamma]) = \sigma_f([\gamma])$ vanishes.

Next, we consider the case where f is a co-orientable flat front. The orientability of M^2 has already been proved in [21]. We set (see §3 of [20])

$$|\rho_\delta| := e^{-2\delta} |\theta/dh| = e^{-2\delta} |Q/dh^2|.$$

Then the singular set of the parallel front f_δ is given by the 1-level set $|\rho_\delta| = 1$. We fix an element $[\gamma] \in \pi_1(M^2)$, where γ is a loop in M^2 . Then we can choose γ so that it does not pass through the zeros of dh nor of θ . Then there exists a constant $c > 0$ such that $|Q/dh^2| \geq c$ on γ . If we take δ sufficiently small, then we have $e^{-2\delta} |Q/dh^2| > 1$ on γ . Then f_δ has no singular point on γ . Thus $\sigma_{f_\delta}([\gamma]) = \sigma_f([\gamma])$ vanishes. $\square$

Remark 3.10. In contrast to the case of H^3 , there are many flat fronts in $\mathbf{R}^3$ with non-trivial zig-zag representations. For example, consider a cylinder over the planar curve in Figure 1.

Remark 3.11. An immersion $f : M^2 \rightarrow \mathbf{R}^3$ is called a *linear Weingarten surface of minimal type* if the mean curvature H and the Gaussian curvature K satisfy the relation $H = aK$, where a is a real constant. Such surfaces in $\mathbf{R}^3$ are the corresponding analogue of linear Weingarten surfaces of Bryant type. As in the method of Section 1, one can define *linear Weingarten fronts of minimal type*. One can easily prove that each linear Weingarten front $f : M^2 \rightarrow \mathbf{R}^3$ of minimal type contains a minimal surface $f_0 : M^2 \rightarrow \mathbf{R}^3$ as a wave front in its parallel family. Since minimal surfaces have a well-known Weierstrass representation formula, the first fundamental form of f_0 is given by $|Q|^2/d\sigma^2$, where Q is the Hopf differential and $d\sigma^2 := 4|dg|^2/(1+|g|^2)^2$ for the Gauss map g of f (and f_0). Since the singular points of f_0 are isolated, the orientability of M^2 is equivalent to the co-orientability of f . (This is a different phenomenon from the case of linear Weingarten fronts of Bryant type. In fact, non-orientable minimal immersions are known.) By the same argument as in the above proof, the zig-zag representation of a given linear Weingarten front of minimal type $f : M^2 \rightarrow \mathbf{R}^3$ vanishes if f is co-orientable.

3.3. An explicit formula for G_* . Recall that the hyperbolic Gauss map of a co-orientable front f is given by $G = [f + \nu]$. On the other hand, we set $G_* := [f - \nu]$, which is also called the (*opposite*) *hyperbolic Gauss map*. For a co-orientable flat front, it is known that G_* is also a meromorphic function, as well as G (cf. [13]). We give here an explicit formula for G_* :

Proposition 3.12. *Under the notation of Theorem 3.1, the opposite hyperbolic Gauss map G_* is given by*

$$(3.9) \quad G_* = G - \frac{(G_h)^2(1 + \varepsilon|h|^2)}{\varepsilon \bar{h} G_h + (G_{hh}/2)(1 + \varepsilon|h|^2)}.$$

Proof. The matrices $\mathcal{A}$ and $\mathcal{B}$ in Theorem 3.1 split as

$$\mathcal{A} = \Phi \Phi^*, \quad \mathcal{B} = \Phi e_3 \Phi^*,$$

where $\Phi^* := {}^t \bar{\Phi}$ and

$$\Phi := \frac{i}{\sqrt{1 + \varepsilon|h|^2}} \begin{pmatrix} -1 & -\varepsilon \bar{h} \\ 0 & 1 + \varepsilon|h|^2 \end{pmatrix}, \quad e_3 := \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Hence we obtain

$$f - \nu = \mathcal{G}(\mathcal{A} - \mathcal{B})\mathcal{G}^* = 2\mathcal{G}\Phi \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} (\mathcal{G}\Phi)^*.$$

Setting $\mathcal{G}\Phi = \begin{pmatrix} p & q \\ r & s \end{pmatrix}$, we have $f - \nu = 2 \begin{pmatrix} q \\ s \end{pmatrix} \begin{pmatrix} \bar{q} & \bar{s} \end{pmatrix}$. This implies $G_* = [f - \nu] = q/s \in \mathbf{C} \cup \{\infty\}$. Then by a straightforward calculation, we get the assertion. $\square$

We immediately get the following:




FIGURE 1. A planar curve of zig-zag number 3

Corollary 3.13. G_* is meromorphic, as well as G , if and only if $\varepsilon = 0$.

Let D be the diagonal set of $S^2 \times S^2$. Then the space $L(H^3)$ of oriented geodesics of hyperbolic 3-space H^3 can be identified with $S^2 \times S^2 \setminus D$ which is a parahermitian symmetric space $SO^0(1, 3)/(SO(2) \cdot \mathbf{R}^*)$ (see Kanai [17], Kaneyuki [18] and also [6]). Recently, Georgiou and Guilfoyle [12] proved that $L(H^3)$ has a canonical neutral Kähler structure and the pair of hyperbolic Gauss maps

$$(G, G_*) : M^2 \rightarrow S^2 \times S^2 \setminus D$$

gives a Lagrangian surface with zero Gaussian curvature if f is a Weingarten surface. Since our explicit formula (3.9) for G_* is written in terms of only G , h and their derivatives, it might be useful for constructing such surfaces with positive genus.

4. ORIENTABILITY AND CO-ORIENTABILITY OF CMC-1 FACES IN S_1^3

In this section, we shall give an application of our explicit formula (3.9) for G_* . Let M^2 be a Riemann surface. A holomorphic map $F : M^2 \rightarrow SL(2, \mathbf{C})$ is called *null* if the determinant of the derivative dF/dz with respect to each local complex coordinate z vanishes identically. It is well known that CMC-1 surfaces in H^3 and in S_1^3 are both projections of null holomorphic curves $F : \tilde{M}^2 \rightarrow SL(2, \mathbf{C})$, where $\tilde{M}^2$ is the universal covering of M^2 and $H^3 = SL(2, \mathbf{C})/SU(2)$ and $S_1^3 = SL(2, \mathbf{C})/SU(1, 1)$. Suppose that a CMC-1 surface in S_1^3 is given by $f = -\mathcal{GBG}^*$ for $\varepsilon = -1$ as in Theorem 3.1. (Here we suppose $f = -\mathcal{GBG}^*$ instead of $\mathcal{GBG}^*$ because it makes the argument below more compatible to [7] and [9]. $-f$ and f differ merely by congruence in S_1^3 , that is, they are essentially the same.) Then the null holomorphic lift F of f is given by

$$(4.1) \quad F = \mathcal{G} \begin{pmatrix} 0 & -i \\ -i & ih \end{pmatrix}$$

Conversely, the projection $Fe_3F^* : M^2 \rightarrow S_1^3$ may admit singular points. Such a surface is called a CMC-1 face. More precisely, a C^∞ map $f : M^2 \rightarrow S_1^3$ is called a *CMC-1 face* if the regular set is open dense in M^2 , and for each $p \in M^2$, there exist a neighborhood U of p and a null holomorphic immersion $F : U \rightarrow SL(2, \mathbf{C})$ such that it has an expression $f = Fe_3F^*$ on U (see [7] and [9]). Note that we usually take a null holomorphic lift F defined on the universal cover $\tilde{M}^2$ for a CMC-1 face.

We shall show in this section the orientability and the co-orientability of CMC-1 faces in this setting. To show the orientability, we will need the explicit formula for G_* , which was a formula missing in [9]. It should also be remarked that CMC-1 faces are not fronts in general but are all frontals, as seen below (see (4.4)). On the other hand, a CMC-1 front in S_1^3 may admit isolated singular points in M^2 , as seen in Corollary 3.4, although CMC-1 faces do not. This implies that the set of CMC-1 fronts is not included in the set of CMC-1 faces and vice versa.

For a CMC-1 face $f = Fe_3F^* (= -\mathcal{GBG}^*)$, we have the following:

$$(4.2) \quad F^{-1}dF = \begin{pmatrix} h & -h^2 \\ 1 & -h \end{pmatrix} \frac{Q}{dh}, \quad dFF^{-1} = \begin{pmatrix} G & -G^2 \\ 1 & -G \end{pmatrix} \frac{Q}{dG},$$

where G , h and Q are corresponding to the data in Theorem 3.1. The formulas (4.2) follow from (3.7) and (4.1). Note that the Hopf differential Q and the secondary Gauss map g have already been introduced in [7]. In our notation, the Hopf differential is denoted by the same Q , the secondary Gauss map g coincides with h , and $\omega := Q/dg$ in [7] is equal to $\theta = Q/dh$.

The singular set S_f of a CMC-1 face $f = Fe_3F = -\mathcal{GBG}^*$ is given by

$$S_f = \{p \in M^2; |h(p)| = 1\}.$$

(See [7, Theorem 1.9].) Away from the singular set S_f , we can define a unit normal field $\nu : M^2 \setminus S_f \rightarrow H_+^3 \cup H_-^3$ by $\nu = \mathcal{GAG}^*$. It follows from (4.1) that

$$(4.3) \quad \nu = F \begin{pmatrix} ih & i \\ i & 0 \end{pmatrix} \mathcal{A} \begin{pmatrix} ih & i \\ i & 0 \end{pmatrix}^* F^* = \frac{\tilde{\nu}}{1 - |h|^2},$$

where

$$(4.4) \quad \tilde{\nu} := F \begin{pmatrix} 1 + |h|^2 & 2h \\ 2\bar{h} & 1 + |h|^2 \end{pmatrix} F^*.$$

(See also [7, Remark 1.2].) Since the equivalence class $[\tilde{\nu}]$ gives a section of M^2 into the projective tangent bundle $P(TS_1^3)$, an arbitrary CMC-1 face f is a frontal. (The definition of a frontal is given in the introduction.) Generic singular points on CMC-1 faces are cuspidal edges, swallowtails and cuspidal cross caps (see [10]). However, cuspidal cross caps never appear on fronts. Hence, CMC-1 faces are not fronts in general.

Theorem 4.1. *Let $f : M^2 \rightarrow S_1^3$ be a CMC-1 face. Then M^2 is orientable.*

Proof. Suppose, by way of contradiction, that M^2 is not orientable. There exists a double covering $\hat{\pi} : \hat{M}^2 \rightarrow M^2$ such that $\hat{M}^2$ is orientable. Set $\hat{f} := f \circ \hat{\pi}$. Let $T : \hat{M}^2 \rightarrow \hat{M}^2$ denote the covering involution which is orientation-reversing. We fix a regular point $p \in \hat{M}^2$. Then there exists a simply connected neighborhood $U \subset \hat{M}^2$ of p such that the restriction $\hat{f}|_U$ is an immersion. Let $\hat{\nu}$ be the unit normal field on U , and $\hat{G}, \hat{G}_*$ denote the hyperbolic Gauss maps for $\hat{f}$ on U . Then $\hat{f} \circ T = \hat{f}$, and either $\hat{\nu} \circ T(q) = \hat{\nu}(q)$ or $\hat{\nu} \circ T(q) = -\hat{\nu}(q)$ holds for each $q \in U$ as vectors in $\mathbf{R}_1^4$. First, we suppose $\hat{\nu} \circ T = -\hat{\nu}$. Then

$$\hat{G} \circ T = [\hat{f} \circ T + \hat{\nu} \circ T] = [\hat{f} - \hat{\nu}] = \hat{G}_*$$

holds on U . This implies that $\hat{G}_*$ is anti-holomorphic, which contradicts Corollary 3.13. Next, we suppose $\hat{\nu} \circ T = \hat{\nu}$. Then, we have

$$\hat{G} \circ T = [\hat{f} \circ T + \hat{\nu} \circ T] = [\hat{f} + \hat{\nu}] = \hat{G},$$

which contradicts that T is orientation-reversing. $\square$

Since G is a holomorphic map, when the unit normal field ν of a CMC-1 face f crosses a singular curve $S_f = \{|h| = 1\}$, the image of ν moves into another sheet of the hyperboloid $H_+^3 \cup H_-^3$. Thus, it is natural to expect ν to be smooth at the singular set under a certain compactification of $H_+^3 \cup H_-^3$. The *hyperbolic 3-sphere* $\bar{\mathcal{H}}^3$ is a 3-dimensional manifold diffeomorphic to the 3-sphere

$$\bar{\mathcal{H}}^3 := \mathbf{R}^3 \cup \{\infty\} \cong S^3$$

endowed with the metric $4dx \cdot dx / (1 - |x|^2)^2$ on $S^3 \setminus \{\text{equator}\}$, where $x := (x_1, x_2, x_3) \in \mathbf{R}^3 \cup \{\infty\}$. We consider the stereographic projection

$$(4.5) \quad \varphi : H_+^3 \cup H_-^3 \ni (x_0, x_1, x_2, x_3) \mapsto \frac{(x_1, x_2, x_3)}{1 - x_0} \in \bar{\mathcal{H}}^3,$$

which is an isometric embedding, and $\bar{\mathcal{H}}^3$ can be considered as a compactification of $H_+^3 \cup H_-^3$.

Theorem 4.2. *The unit normal field of a CMC-1 face can be considered as a real analytic map into the hyperbolic 3-sphere $\bar{\mathcal{H}}^3$. In other words, the map $\varphi \circ \nu$, which is defined on $M^2 \setminus S_f$, can be real analytically extended across the singular set S_f .*

Proof. We set $F = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$. Then A, B, C, D are holomorphic functions that are locally defined on a coordinate neighborhood of M^2 . We have only to show that $\varphi \circ \nu$ is real analytic at the point p where $|h(p)| = 1$, since $S_f = \{p \in M^2; |h(p)| = 1\}$.

By a straightforward calculation using (3.6), (4.3), (4.4) and (4.5), one can check that each component of $\varphi \circ \nu$ is a rational expression in A, B, C, D, h and their conjugates, whose denominator r is the following:

$$r = 2(1 - |h|^2) + |A + B\bar{h}|^2 + |C + D\bar{h}|^2 + |Ah + B|^2 + |Ch + D|^2.$$

Hence, it is sufficient to show that $r(p) \neq 0$ when $|h(p)| = 1$. However, the equality

$$Ah + B = A + B\bar{h} = C + D\bar{h} = Ch + D = 0$$

holds at p only when $F(p)$ is a singular matrix. Therefore $r(p)$ cannot be zero. This proves the assertion. $\square$

Corollary 4.3. *Let $f : M^2 \rightarrow S_1^3$ be a CMC-1 face. Then there exists a real analytic normal vector field defined on M^2 . In particular, f is a co-orientable frontal.*

Proof. Define a map $\psi : H_+^3 \cup H_-^3 \rightarrow \mathbf{R}_1^4$ by

$$\psi(u) = \begin{cases} u/|u|_E & \text{if } u \in H_+^3, \\ -u/|u|_E & \text{if } u \in H_-^3, \end{cases}$$

where $|u|_E$ denotes the Euclidean norm of u as a vector in $\mathbf{R}^4 (= \mathbf{R}_1^4)$. For the unit normal field $\nu : M^2 \setminus S_f \rightarrow H_+^3 \cup H_-^3$, the equality $\psi \circ \nu = \psi \circ \varphi^{-1} \circ N$ holds on $M^2 \setminus S_f$. Here, $N : M^2 \rightarrow \bar{\mathcal{H}}^3$ denotes the real analytic extension of $\varphi \circ \nu : M^2 \setminus S_f \rightarrow \bar{\mathcal{H}}^3$ explained in Theorem 4.2. On the other hand, we can calculate $\psi \circ \varphi^{-1} : \bar{\mathcal{H}}^3 \setminus \{|x| = 1\} \rightarrow \mathbf{R}_1^4$ as follows:

$$\psi \circ \varphi^{-1} : (x, y, z) \mapsto \frac{1}{\sqrt{\Delta}}(1 + x^2 + y^2 + z^2, -2x, -2y, -2z)$$

where

$$\Delta := (x^2 + y^2 + z^2 + 1)^2 + 4x^2 + 4y^2 + 4z^2.$$

It follows that $\psi \circ \varphi^{-1}$ can be smoothly extended to the whole of $\bar{\mathcal{H}}^3$. Therefore $\Psi := \psi \circ \varphi^{-1} \circ N$ can be considered as a smooth map on M^2 such that $\Psi(p) \in \mathbf{R}_1^4$ is perpendicular to the position vector $f(p) \in \mathbf{R}_1^4$ and the subspace $df(T_p M) (\subset \mathbf{R}_1^4)$ at every regular point $p \in M^2 \setminus S_f$. Thus Ψ gives a globally defined normal vector field of f , which proves the co-orientability of f . $\square$

5. CO-ORIENTABILITY OF MAXIMAL SURFACES IN $\mathbf{R}_1^3$

In this section, we shall discuss the co-orientability of maximal surfaces in Lorentz-Minkowski 3-space given as a joint work with Shoichi Fujimori and Kotaro Yamada during the authors' stay at RIMS in Kyoto and a meeting of the second author with Kotaro Yamada at Osaka University on June 2009: CMC-1 faces in S_1^3 have very similar properties to maximal surfaces in Lorentz-Minkowski 3-space $\mathbf{R}_1^3$

of signature $(-, +, +)$. In [28], maxfaces (spacelike maximal surfaces with admissible singular points) are defined as C^∞ maps of orientable 2-manifolds into $\mathbf{R}_1^3$. If M^2 is non-orientable, we can take a double covering $\hat{\pi} : \hat{M}^2 \rightarrow M^2$ such that $\hat{M}^2$ is orientable. For a non-orientable 2-manifold M^2 , a C^∞ map $f : M^2 \rightarrow \mathbf{R}_1^3$ is called a *maxface* if $f \circ \pi$ is a maxface in the sense of [28].

In contrast to CMC-1 faces in S_1^3 , there are non-orientable complete maxfaces with or without handles (see Fujimori and López [8]). Figure 2 shows two different non-orientable maxfaces given in [8]. However, orientability and co-orientability of maxfaces are not discussed explicitly in [28].




FIGURE 2. Non-orientable maxfaces given in [8] (the picture is produced by Fujimori)

The *hyperbolic 2-sphere* $\bar{\mathcal{H}}^2$ is a Riemann sphere

$$\bar{\mathcal{H}}^2 := \mathbf{C} \cup \{\infty\} \cong S^2$$

endowed with the metric $4|dw|^2/(1-|w|^2)^2$ on $S^2 \setminus \{\text{equator}\}$, where $w \in \mathbf{C} \cup \{\infty\}$. The hyperboloid of two sheets in $\mathbf{R}_1^3$, denoted by $H_+^2 \cup H_-^2$, is

$$H_+^2 \cup H_-^2 := \{(x_0, x_1, x_2) \in \mathbf{R}_1^3; -x_0^2 + x_1^2 + x_2^2 = -1\}.$$

We consider the stereographic projection

$$\varphi : H_+^2 \cup H_-^2 \ni (x_0, x_1, x_2) \mapsto \frac{x_1 + ix_2}{1 - x_0} \in \bar{\mathcal{H}}^2,$$

which is an isometric embedding, and $\bar{\mathcal{H}}^2$ can be considered as a compactification of $H_+^2 \cup H_-^2$. The following fact is well-known (cf. [28]):

Fact 5.1. *Let M^2 be a Riemann surface, $f : M^2 \rightarrow \mathbf{R}_1^3$ a maxface, and $G : M^2 \rightarrow \bar{\mathcal{H}}^2 = \mathbf{C} \cup \{\infty\}$ its Lorentzian Gauss map as a meromorphic function. Then $\nu = \varphi^{-1} \circ G$ gives a unit normal vector field defined on the regular set of f .*

Using this, we shall prove the following assertion regardless of whether M^2 is orientable or not.

Proposition 5.2. ([11]) *Let $f : M^2 \rightarrow \mathbf{R}_1^3$ be a maxface. Then there exists a real analytic normal vector field defined on M^2 . In particular, f is a co-orientable frontal.*

Proof. A smooth map $F : M^2 \rightarrow \mathbf{R}_1^3$ is a co-orientable frontal if and only if there exists a C^∞ vector field η defined on M^2 such that $\eta_p \in T_{F(p)}\mathbf{R}_1^3$ is Lorentzian orthogonal to the vector space $dF(T_p M^2)$.

We define a map $\psi : H_+^2 \cup H_-^2 \rightarrow \mathbf{R}_1^3$ by

$$\psi(u) = \begin{cases} u/|u|_E & \text{if } u \in H_+^2, \\ -u/|u|_E & \text{if } u \in H_-^2, \end{cases}$$

where $|u|_E$ denotes the Euclidean norm of u as a vector in $\mathbf{R}^3 (= \mathbf{R}_1^3)$.

Suppose that M^2 is orientable. By Fact 5.1,

$$(5.1) \quad \nu := \psi \circ \varphi^{-1} \circ G$$

gives a Lorentzian normal vector field of f on $M^2 \setminus S_f$. On the other hand, we can calculate $\psi \circ \varphi^{-1} : \tilde{\mathcal{H}}^2 \setminus \{|w| = 1\} \rightarrow \mathbf{R}_1^3$ as follows:

$$\psi \circ \varphi^{-1} : w = x + iy \mapsto \frac{1}{\sqrt{\Delta}}(1 + x^2 + y^2, -2x, -2y),$$

where

$$\Delta := (x^2 + y^2 + 1)^2 + 4x^2 + 4y^2.$$

It follows that $\psi \circ \varphi^{-1}$ can be smoothly extended to the whole of $\tilde{\mathcal{H}}^2$. Therefore ν given in (5.1) is a globally defined normal vector field of f . In fact, by a straightforward calculation, we have that

$$(5.2) \quad \nu = \frac{1}{\sqrt{(1 + |G|^2)^2 + 4|G|^2}}(1 + |G|^2, -2\operatorname{Re}(G), -2\operatorname{Im}(G)).$$

So the orientability implies the co-orientability. Next, suppose that $f : M^2 \rightarrow \mathbf{R}_1^3$ is a maxface and M^2 is non-orientable. Then there exists a double covering $\hat{\pi} : \hat{M}^2 \rightarrow M^2$ such that $\hat{M}^2$ is orientable. Set $\hat{f} := f \circ \hat{\pi}$ and denote by $G : M^2 \rightarrow \mathbf{C} \cup \{\infty\}$ the Gauss map of the maxface $\hat{f}$. Let $T : \hat{M}^2 \rightarrow \hat{M}^2$ be the covering involution of the double covering $\hat{\pi} : \hat{M}^2 \rightarrow M^2$. Then $(\varphi^{-1} \circ G) \circ T(p)$ is antipodal to $(\varphi^{-1} \circ G)(p)$ in $H_+^2 \cup H_-^2$ for every $p \in \hat{M}^2$ (cf. [8]). It follows that

$$(5.3) \quad G \circ T(p) = 1/\overline{G(p)} \quad (p \in \hat{M}^2).$$

By (5.3), the vector field ν given by (5.2) is a Lorentzian normal vector field on $\hat{M}^2$ satisfying $\nu \circ T = \nu$. In other words, ν can be considered as a smooth normal vector field on M^2 , which proves the assertion. $\square$

Remark 5.3. Let $f : M^2 \rightarrow \mathbf{R}_1^3$ be a non-orientable maxface. Then a closed regular curve $\gamma : [0, 1] \rightarrow M^2$ is called a *non-orientable loop* if the orientation of M^2 reverses along the curve. Then as pointed out in Fujimori and López [8], there is at least one singular point on γ . Take a double covering $\hat{\pi} : \hat{M}^2 \rightarrow M^2$ such that $\hat{M}^2$ is orientable. Set $\hat{f} := f \circ \hat{\pi}$ and denote by $G : \hat{M}^2 \rightarrow \tilde{\mathcal{H}}^2 = \mathbf{C} \cup \{\infty\}$ the Lorentzian Gauss map of the maxface $\hat{f}$. Since the singular set $S_{\hat{f}}$ of $\hat{f}$ is the 1-level set of $|G|$, it consists of a union of regular curves outside of the discrete set

$$\Gamma := \left\{ p \in S_{\hat{f}}; dG(p) = 0 \right\}.$$

Here G is considered as a smooth map into S^2 and dG is the derivative of G . A non-orientable loop γ is called *generic* if its lift $\hat{\gamma}$ on $\hat{M}^2$ meets the singular set only at finitely many points on $S_{\hat{f}} \setminus \Gamma$, where $\hat{\gamma}$ passes through these points transversally to the set $S_{\hat{f}} \setminus \Gamma$. We fix a generic non-orientable loop γ arbitrarily. Since $T \circ \hat{\gamma}(0) = \hat{\gamma}(1)$, by (5.3), we have $|G \circ \hat{\gamma}(0)| = 1/|G \circ \hat{\gamma}(1)|$. Since the curve $G \circ \hat{\gamma}(t)$ ($t \in [0, 1]$) is continuous, it meets the circle $S^1 = \{w \in \mathbf{C} \cup \{\infty\}; |w| = 1\}$

odd times. Thus *the number of singular points on a generic non-orientable loop γ is odd.* (In fact, the number of connected components of the singular set on the non-orientable maxfaces in Figure 2 is one and three, respectively.)

APPENDIX A. DEGENERATE WEINGARTEN FRONTS

In hyperbolic 3-space H^3 , the following facts are well-known;

- (1) one of the parallel surfaces of an open portion of a hyperbolic cylinder degenerates to a hyperbolic line, and
- (2) one of the parallel surfaces of an open portion of a geodesic sphere (a compact totally umbilical surface) degenerates to a single point.

Note that these are Weingarten fronts. (A *Weingarten front* is defined by Definition 2.5 removing the adjective ‘horospherical linear’. This definition makes sense based on the fact that any parallel surface of a Weingarten surface is again Weingarten.) The above fronts (1) and (2) are called a *degenerate cylinder* and a *degenerate sphere*, respectively. We wish to prove a converse statement. Before this, let us recall the following well known lemma:

Lemma A.1. *Let $f : M^2 \rightarrow H^3$ be an immersion, and κ_i ($i = 1, 2$) the principal curvatures. Then the set of singular points of a parallel surface f_δ coincides with $\{p \in M^2 ; \coth^{-1} \kappa_i(p) = \delta\}$.*

Lemma A.1 asserts that $p \in M^2$ is a singular point of f_δ if and only if δ is the radius $\coth^{-1} \kappa_i(p)$ of the principal curvature $\kappa_i(p)$. It is easily observed that any parallel surface is free from singular points if and only if $|\kappa_i| \leq 1$.

Proposition A.2. *Let $f : M^2 \rightarrow H^3$ be a Weingarten front. Then all points in M^2 are singular points if and only if f is locally a degenerate cylinder or a degenerate sphere.*

Proof. Suppose that all points in M^2 are singular points of f . Let $p \in M^2$ be an arbitrary point. There exist a neighborhood U of p and a real number δ_0 such that $f_{\delta_0} : U \rightarrow H^3$ is an immersion. In other words, $f : U \rightarrow H^3$ is a parallel surface of an immersion $g := f_{\delta_0}$, i.e., $f = g_{-\delta_0}$. Since all points are singular, it follows from Lemma A.1 that

$$-\delta_0 = \coth^{-1} \kappa(p) \quad \text{for all } p \in U,$$

where κ denotes one of the principal curvatures of g . It implies that κ is constant on U and satisfies $|\kappa| > 1$. Moreover, another principal curvature is also constant because g is Weingarten. Thus g is an isoparametric surface. By the classification due to Cartan [4], the immersion g is totally umbilical or locally a standard embedding of $S^1 \times H^1$ into H^3 . Under our assumption, the former is a part of geodesic sphere because $|\kappa| > 1$, the latter is a part of a hyperbolic cylinder.

Finally, by the connectivity of M^2 and by the continuity of f , we can conclude that $f : M^2 \rightarrow H^3$ is a degenerate cylinder or a degenerate sphere. $\square$

Acknowledgement The authors thank Antonio Martínez for a fruitful discussion on the geometry of linear Weingarten surfaces, when they visited the Fukuoka conference on November, 2004. They also thank Shyuichi Izumiya for his informal lecture on the duality between fronts in H^3 and S_1^3 at the Karatsu workshop on October, 2005. The first author thanks Shoichi Fujimori for fruitful discussions about the proof of Proposition 3.12, after the Karatsu workshop. The second

author thanks Masahiko Kanai, who pointed out the importance of $L(H^3)$ as the range of hyperbolic Gauss maps of CMC-1 surfaces in H^3 when the second author stayed at Nagoya University in 2001. Finally, the authors thank Kotaro Yamada, Wayne Rossman and Francisco López for valuable comments.

REFERENCES

- [1] J. Aledo and J. Espinar, A conformal representation for linear Weingarten surfaces in the de Sitter space, *J. Geom. Phys.* **57**, 1669–1677 (2007).
- [2] J. Aledo and J. A. Galvez, Complete surfaces in hyperbolic space with a constant principal curvature, *Math. Nachr.* **278**, 1111–1116 (2005).
- [3] R. Bryant, Surfaces of mean curvature one in hyperbolic space, *Astérisque* **154–155**, 321–347 (1987).
- [4] É. Cartan, Familles de surfaces isoparamétriques dans les espaces à courbure constante, *Ann. di Mat.*, **17**, 177–191 (1938).
- [5] B. Y. Chen, Minimal surfaces with constant Gauss curvature, *Proc. Amer. Math. Soc.* **34**, 504–508 (1972).
- [6] V. Cruceanu, P. Fortuney and P. M. Gadea, A survey on paracomplex geometry, *Rocky Mountain J. Math.* **26**, 83–115 (1996).
- [7] S. Fujimori, Spacelike CMC 1 surfaces with elliptic ends in de Sitter 3-Space, *Hokkaido Math. J.* **35**, 289–320 (2006).
- [8] S. Fujimori and F. J. López, Nonorientable maximal surfaces in the Lorentz-Minkowski 3-space, preprint, arXiv:0905.2113.
- [9] S. Fujimori, W. Rossman, M. Umehara, K. Yamada and S.-D. Yang, Spacelike mean curvature one surfaces in de Sitter 3-space, *Comm. Anal. Geom.*, **17**, 383–427 (2009) .
- [10] S. Fujimori, K. Saji, M. Umehara and K. Yamada, Singularities of maximal surfaces, *Math. Z.* **259**, 827–848 (2008).
- [11] S. Fujimori, M. Kokubu, M. Umehara and K. Yamada, Personal meetings at RIMS Kyoto and at Osaka University on June, 2009.
- [12] N. Georgiou and B. Guilfoyle, A characterization of Weingarten surfaces in hyperbolic 3-space, preprint, arXiv:0709.2441.
- [13] J. A. Gálvez, A. Martínez and F. Milán, Flat surfaces in hyperbolic 3-space, *Math. Ann.* **316**, 419–435 (2000).
- [14] J. A. Gálvez, A. Martínez and F. Milán, Complete linear Weingarten surfaces of Bryant type. A Plateau problem at infinity, *Trans. Amer. Math. Soc.* **356**, 3405–3428 (2004).
- [15] S. Izumiya, D. Pei, M. C. Romero-Fuster and M. Takahashi, The horospherical geometry of submanifolds in Hyperbolic space, preprint.
- [16] S. Izumiya, K. Saji and M. Takahashi, Horospherical flat surfaces in Hyperbolic 3-space, preprint.
- [17] M. Kanai, Geodesic flows of negatively curved manifolds with smooth stable and unstable foliations, *Ergodic Theory Dynam. Systems* **8**, 215–239 (1988).
- [18] S. Kaneyuki and M. Kozai, Paracomplex structures and affine symmetric spaces, *Tokyo J. Math.* **8**, 81–98 (1985).
- [19] M. Kokubu, Surfaces and fronts with harmonic-mean curvature one in hyperbolic three-space, *Tokyo J. Math.* **32**, 177–200 (2009).
- [20] M. Kokubu, W. Rossman, K. Saji, M. Umehara and K. Yamada, Singularities of flat fronts in hyperbolic space, *Pacific J. Math.* **221**, 303–351 (2005).
- [21] M. Kokubu, W. Rossman, M. Umehara and K. Yamada, Flat fronts in hyperbolic 3-space and their caustics, *J. Math. Soc. Japan* **59**, 265–299 (2007).
- [22] M. Kokubu, M. Umehara and K. Yamada, Flat fronts in hyperbolic 3-space, *Pacific J. Math.* **216**, 149–175 (2004).
- [23] R. Langevin, G. Levitt and H. Rosenberg, Classes d’homotopie de surfaces avec rebroussements et queues d’aronde dans $\mathbf{R}^3$, *Canad. J. Math.* **47**, 544–572 (1995).
- [24] K. Saji, M. Umehara and K. Yamada, The geometry of fronts, *Ann. of Math.* **169**, 491–529 (2009).
- [25] K. Saji, M. Umehara and K. Yamada, A_k singularities of wave fronts, to appear in *Math. Proc. Cambridge Philos. Soc.* **146**, 731–746 (2009).

- [26] C. Takizawa and K. Tsukada, Horocyclic surfaces in Hyperbolic 3-space, Kyusyu J. Marh. **63**, 269–284 (2009).
- [27] M. Umehara and K. Yamada, Complete surfaces of constant mean curvature-1 in the hyperbolic 3-space, Ann. of Math. **137**, 611–638 (1993).
- [28] ———, Maximal surfaces with singularities in Minkowski space, Hokkaido Math. J. **35**, 13–40 (2006).

(Kokubu) DEPARTMENT OF MATHEMATICS, SCHOOL OF ENGINEERING, TOKYO DENKI UNIVERSITY, CHIYODA-KU, TOKYO 101-8457, JAPAN

E-mail address: kokubu@cck.dendai.ac.jp

(Umehara) DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

E-mail address: umehara@math.sci.osaka-u.ac.jp